

Lymphoedema Therapy Referral

Patient Information or Bradma

Name: _____ NHI: _____
 Address: _____

 Phone: _____ Mobile: _____

Date: _____ Diagnosis: _____

Relevant Information: Include history of disease, spec pathology causing high protein lymphoedema, sites involved, patient status and prognosis. Blood test/medications etc.

The Management of Lymphoedema in Advanced Cancer and Oedema at the End of Life - International Guidelines see: www.lympho.org

Patient profile: - please tick relevant boxes (See over for lymphoedema treatment indications and limitations)

- Lives within the Nelson-Richmond city boundaries
- Lives alone or with frail spouse
- Lives with able bodied carer or receives care from home support agency
- Has good limb function and is mobile
- Is inactive
- Pain is present
- Can understand and participate in treatment
- Can attend the hospice
- Has severe cardiac or renal failure
- Could tolerate bandaging or compression garments
- Has skin leakage
- Has peripheral vascular disease or diabetes
- Has history of DVT

Referred by: _____ Designation: _____

Authorised by: _____ Dr/HMO/PC Nurse Practitioner _____

Office Use Only Accepted, date _____ Declined, date _____

More info. required: _____ Letter sent, date: _____ Phone to discuss, date: _____

Lymphoedema Treatment Indications and Limitations

Exercise & Elevation

Anytime

Active or Passive

Kinesiotaping

Indications:

- Distinct area of overall swelling
- Scrotal, thigh and buttock oedema
- Patient can attend clinic for taping

Contraindications:

- Sensitive or allergic skin
- Fragile or broken areas
- Receiving end of life care and treatment deemed inappropriate

Massage

Indications:

- Patient can tolerate massage
- Patient can attend Hospice clinic for massage

Contraindications:

- Skin breakdown
- Receiving end of life care and treatment deemed inappropriate

Bandaging

Indications:

- Wishes to increase mobility or functional ability
- Has lower leg and ankle lymphoedema or
- Has lower arm lymphoedema
- Circumferential difference of more than 2 cm between affected and non-affected limb
- Skin leakage developing
- High risk of pressure areas on lower limb
- Lymphoedema nurse determines effectiveness of bandaging

Contraindications:

- Severe cardiac or renal disease
- Immobility
- Patient cannot tolerate bandaging
- Receiving end of life care and treatment deemed inappropriate

Compressions Garments

Indications:

- Patient has a prognosis of 3 months or more
- Useful for limbs only, not body area
- Patient has good limb function and is mobile
- Careers willing to assist with applying garments or Support Care Worker arranged

Contraindications:

- Patient has severe peripheral vascular disease or a condition (e.g. diabetes) causing inadequate arterial blood flow:
- Skin breakdown or serous ooze
- Cardiac failure
- Receiving end of life care and treatment deemed inappropriate.

Adapted from Nurse Maude Hospice, Christchurch